



THE
ANSWER
SERIES *Your Key to Exam Success*

DBE
PAST PAPERS

Maths Toolkit


OFFICIAL EXAMS & MEMOS

Anne Eadie, Gretel Lampe, Jenny Campbell & Susan Carletti

GRADE

12

CAPS



Grade 12 Maths Toolkit | DBE Past Papers

OFFICIAL EXAMS & MEMOS

The Answer Series Grade 12 Maths Toolkit is a low-priced product, offering both theory and practice, and is perfect for exam preparation for matrics.

This **UP-TO-DATE** publication is indeed a TOOLKIT, containing:

DBE Nov Paper 1 & Paper 2 Exams (2016 – 2023) with **comprehensive solutions** to all papers, including TOPIC GUIDES that make it possible to select questions on **separate topics**. **Challenging questions**, aligned with DBE Diagnostic Reports, have been clearly indicated throughout this study guide.

Supportive, vital documents & powerful summaries

- mark distribution and cognitive levels
- the curriculum
- all examinable proofs
- summaries on trigonometry, quadrilaterals, concavity, analytical geometry and circle geometry,
- theorem statements and acceptable reasons
- calculator instructions
- DBE formulae / information sheet

How learners can improve their exam techniques:

- write a few of the papers under exam conditions
- get comfortable with having to concentrate for the full 3 hour time period
- learn to work through the paper a few times, answering all the routine questions first, then coming back for more challenging questions that take more time, and
- finally, when all else is done, tackling the questions that need more time and attention

Good exam technique makes a huge difference to anyone's ability to produce top quality work under pressure and there is no doubt that The Answer Series Grade 12 Maths Toolkit levels the playing fields and ensures that everyone has equal access to success.



Maths Toolkit

OFFICIAL EXAMS & MEMOS

Anne Eadie, Gretel Lampe, Jenny Campbell & Susan Carletti

Other Gr 12 publications available

► **GRADE 12 MATHEMATICS 2-in-1**

- 1 Questions in topics
- 2 Exam papers
- 3 A separate booklet on challenging, Level 3 & 4 questions

Full solutions provided throughout

► **GRADE 12 MATHEMATICS P & A**

10 additional, challenging practice exams & answers

THIS PAST PAPERS TOOLKIT INCLUDES

- **DBE Exam Papers**
- **Comprehensive solutions to all papers – compiled by our authors, not from the official memoranda**
- **Supportive, vital documents & powerful summaries**

eBook available 



CONTENTS: Gr 12 Exams for DBE

The Exam

Sure Route to Success in Matric Maths

Important Advice for Matrics

The Curriculum (CAPS): Overview of Topics

DBE Paper 1 Topic Guide

DBE Paper 2 Topic Guide

| |
|---------------------------|
| DBE November 2016 Paper 1 |
| DBE November 2016 Paper 2 |
| DBE November 2017 Paper 1 |
| DBE November 2017 Paper 2 |
| DBE November 2018 Paper 1 |
| DBE November 2018 Paper 2 |
| DBE November 2019 Paper 1 |
| DBE November 2019 Paper 2 |
| DBE November 2020 Paper 1 |
| DBE November 2020 Paper 2 |
| DBE November 2021 Paper 1 |
| DBE November 2021 Paper 2 |
| DBE November 2022 Paper 1 |
| DBE November 2022 Paper 2 |
| DBE November 2023 Paper 1 |
| DBE November 2023 Paper 2 |

Exam Memo

| | |
|----|-----|
| 1 | |
| 2 | |
| 3 | A1 |
| 5 | A6 |
| 8 | A11 |
| 10 | A15 |
| 13 | A23 |
| 15 | A28 |
| 18 | A32 |
| 20 | A36 |
| 23 | A40 |
| 25 | A45 |
| 28 | A49 |
| 30 | A55 |
| 33 | A61 |
| 36 | A66 |
| 39 | A71 |
| 41 | A77 |

NOTE

The questions marked with an asterisk (*) are identified as challenging . . . average performance < 40%.



VALUABLE DOCUMENTS

| | |
|---|-------|
| Bookwork: Examinable Proofs | i |
| Trig Summary | vii |
| Quadrilaterals – definitions, areas & properties | viii |
| Concavity & The Point of Inflection | ix |
| Analytical Geometry Toolkit | xi |
| Grouping of Circle Geometry Theorems | xiii |
| Euclidean Geometry: Theorem Statements & Acceptable Reasons | xiv |
| Calculator Instructions | xvii |
| DBE Formula/Information Sheet | xviii |



Be sure to incorporate these Theory documents regularly as you revise.



We are grateful to the Department of Basic Education for granting their permission for the inclusion of these exam papers.

| DBE PI: TOPIC GUIDE | 2016 | 2017 | 2018 | 2019 | 2020 | 2021 | 2022 | 2023 |
|--|---|----------------------------------|-----------------------|-----------------------|---------------------------|----------------------|----------------------|----------------------|
| ► Algebra: [25] | | | | | | | | |
| Quadratic equations & theory | 1.1.1, 1.1.2, 1.2.1 | 1.1.1, 1.1.2 | 1.1.1, 1.1.2 | 1.1.1, 1.1.2 | 1.1.1, 1.1.2 | 1.1.1, 1.1.2, 1.3* | 1.1.1, 1.1.2 | 1.1.1, 1.1.2 |
| Quadratic inequalities | 1.2.2 | 1.3.1 | 1.1.3 | 1.1.3 | 1.1.3 | 1.1.3 | 1.1.3 | 1.1.4 |
| Simultaneous equations | 1.3 | 1.2 | 1.2 | 1.2 | 1.2 | 1.2 | 1.2 | 1.2 |
| ► Exponents: | | | | | | | | |
| Expressions | | | 1.3* | | | | 1.3 | |
| Equations & Inequalities | 1.1.4 | | | 1.3* | 1.3* | | 1.4* | 1.3* |
| ► Surds: | | | | | | | | |
| Expressions | | | | | | | | |
| Equations | 1.1.3 | 1.1.3 | 1.1.4 | 1.1.4 | 1.1.4 | 1.1.4 | 1.1.4 | 1.1.3 |
| ► Logs (Application) | | | | | | | | |
| ► Patterns & Sequences: [25] | | | | | | | | |
| Quadratic | 3.1* | 2.1 | | 2.1 | 2.2 | 3.1, 3.2, 3.3*, 3.4* | 3 | 2.2 |
| Arithmetic | 2.1 – 2.3, 2.4* | 2.2 | | | 2.1 | 4.1, 4.2 | | 2.1 |
| Geometric | 3.2* | | 3.1, 3.2 | 2.2 | 11.3* | 2 | 2.1 | 3.1.1 |
| Σ | | 3* | 3.3, 3.4* | 3.1* | 3.1, 3.2* | 4.3*, 4.4 | 2.2 | 3.1.2 |
| Mixed / General | | | 2.1 – 2.3 | 3.2 | | | | 3.2* |
| ► Finance, Growth & Decay: [15] | | | | | | | | |
| Simple & compound growth & decay | | 6.1 | | 6.1 | 6.2 | 8.1, 8.2 | 6.1 | 6.1.1, 6.2.1 |
| Nominal & Effective interest rates | | | | | | | | 6.1.2 |
| Annuities | 7.1 – 7.3, 7.4* | 6.2* | 7.2 | 6.2 | 6.1, 6.3* | 8.3* | 6.2, 6.3* | 6.2.2, 6.3* |
| Time line | | | 7.1* | | | | | |
| ► Functions & Graphs: [35] | | | | | | | | |
| Straight line and/or parabola | | 1.3*, 4.1 – 4.4, 4.5*, 4.6, 4.7* | 6.1 – 6.3, 6.4*, 6.6* | | | 7 | | |
| Hyperbola | | | 5.1 – 5.3, | | 4.1 | 5 | 4.1 | |
| Exponent. & log function (incl. Inverses) | 4.1 – 4.4, 4.5* | | | | | 6.1, 6.2*, 6.3, 6.4* | | |
| Inverse functions | | | 4.1 – 4.3, 4.4* | 5.1 – 5.3, 5.4*, 5.5* | 5.1, 5.2, 5.3*, 5.4*, 5.5 | | 5.1 – 5.4, 5.5* | |
| Mixed | 5.1, 5.2*, 5.3, 5.4*, 5.5*, 6.1, 6.2, 6.3*, 6.4 | 5.1 – 5.5, 5.6* | | 4.1 – 4.6, 4.7* | 4.2 | | 4.2 | 4.1 – 4.7 (4.6*), 5* |
| ► Differential Calculus: [35] | | | | | | | | |
| Finding the derivative: 1 st principles | 8.1, 8.2* | 7.1 | 8.1 | 7.1 | 7.1 | 9.1 | 7.1 | 7.1 |
| Finding the derivative: using the rules | 8.3 | 7.2 | 8.2 | 7.2, 7.3 | 7.2, 8.4 | 9.2 | 7.2 | 7.2 |
| Finding the average gradient | | | | | | | | |
| Tangent: the gradient & the equation | 8.4* | | | 7.4* | | | | 7.3* |
| Curve sketching & f'' & concavity | 5.6*, 9.1, (9.2 – 9.4)* | 8* | 5.4*, 6.5*, 9.1*, 9.2 | 9.1, 9.2*, 9.3, 9.4* | 8.1, 8.2, 8.3* | (10.1 – 10.4)* | 7.3*, 8.1, 8.2, 8.3* | 8.1 – 8.5, (8.3)* |
| Practical application (incl. max/min) | 1.2.3, 10.1, 10.2*, 10.3* | 9* | 10* | 8.1, 8.2*, 8.3 | 8.5*, 9.1*, 9.2* | 11* | 9* | 9* |
| ► Probability: [15] | | | | | | | | |
| Probability rules | | | 12.1 | | | 12.1* | | 10.1 |
| Venn diagrams | | 10.1, 10.2*, 10.3* | | 11.1* | | | 10.1* | |
| Tree diagrams | | | 12.2* | | 11.1*, 11.2 | | | 10.2* |
| 2-way Contingency tables | 11.1, 11.2*, 11.3 | | | | | | | |
| Fundamental Counting Principle | 12* | 11* | 11* | 10*, 11.2 | 10* | 12.2* | 10.2* | 10.3* |

Questions marked with an asterisk (*) were identified as challenging ... ave performance < 40%

DBE NOVEMBER EXAMS



DBE NOV 2016 PAPER 1

Clearly show ALL calculations, diagrams, graphs, etc. that you have used in determining your answers.

Answers only will NOT necessarily be awarded full marks.

You may use an approved scientific calculator (non-programmable and non-graphical), unless stated otherwise.

If necessary, round off answers to **TWO** decimal places, unless stated otherwise.

► ALGEBRA AND EQUATIONS AND INEQUALITIES [24]

QUESTION 1

Answers on p. A1

1.1 Solve for x :

1.1.1 $x(x - 7) = 0$ (2)

1.1.2 $x^2 - 6x + 2 = 0$ (correct to TWO decimal places) (3)

1.1.3 $\sqrt{x-1} + 1 = x$ (5)

1.1.4 $3^{x+3} - 3^{x+2} = 486$ (4)

1.2 Given: $f(x) = x^2 + 3x - 4$

1.2.1 Solve for x if $f(x) = 0$ (2)

1.2.2 Solve for x if $f(x) < 0$ (2)

1.2.3 Determine the values of x for which $f'(x) \geq 0$ (2)

1.3 Solve for x and y : $x = 2y$ and $x^2 - 5xy = -24$ (4) [24]

► PATTERNS & SEQUENCES [26]

QUESTION 2

Answers on p. A1

Given the finite arithmetic sequence:

$5 ; 1 ; -3 ; \dots ; -83 ; -87$

2.1 Write down the fourth term (T_4) of the sequence. (1)

2.2 Calculate the number of terms in the sequence. (3)

2.3 Calculate the sum of all the negative numbers in the sequence. (3)

2.4* Consider the sequence:

$5 ; 1 ; -3 ; \dots ; -83 ; -87 ; \dots ; -4\ 187$

Determine the number of terms in this sequence that will be exactly divisible by 5. (4) [11]

QUESTION 3

Answers on p. A2


3.1* The first four terms of a quadratic number pattern are

$-1 ; x ; 3 ; x + 8$

3.1.1 Calculate the value(s) of x . (4)

3.1.2 If $x = 0$, determine the position of the first term in the quadratic number pattern for which the sum of the first n first differences will be greater than 250. (4)

3.2* Rectangles of width 1 cm are drawn from the edge of a sheet of paper that is 30 cm long such that there is a 1 cm gap between one rectangle and the next. The length of the first rectangle is 21 cm and the length of each successive rectangle is 85% of the length of the previous rectangle until there are rectangles drawn along the entire length of AD. Each rectangle is shaded.



3.2.1 Calculate the length of the 10th rectangle. (3)

3.2.2 Calculate the percentage of the paper that is shaded. (4) [15]

► FUNCTIONS & GRAPHS [35]


QUESTION 4

Answers on p. A2

Sketched below is the graph of $h(x) = a^x$, $a > 0$.

R is the y-intercept of h .

The points P(2; 9) and Q(b; $\frac{1}{81}$) lie on h .



4.1 Write down the equation of the asymptote of h . (1)

4.2 Write down the coordinates of R. (1)

4.3 Calculate the value of a . (2)

4.4 D is a point such that $DQ \parallel y$ -axis and $DP \parallel x$ -axis. Calculate the length of DP. (4)

4.5* Determine the values of k for which the equation $h(x + 2) + k = 0$ will have a root that is less than -6 . (3) [11]

Finding some challenges along the way?




Use the memo wisely and make sure you process and understand every step of the solution.

QUESTION 4

Answers on p. A17

In the diagram, $P(-4; 5)$ and $K(0; -3)$ are the end points of the diameter of a circle with centre M . S and R are respectively the x - and y -intercepts of the tangent to the circle at P . θ is the inclination of PK with the positive direction of the x -axis.



4.1 Determine:

- 4.1.1 The gradient of SR (4)
- 4.1.2 The equation of SR in the form $y = mx + c$ (2)
- 4.1.3 The equation of the circle in the form $(x - a)^2 + (y - b)^2 = r^2$ (4)
- 4.1.4 The size of \hat{PKR} (3)
- 4.1.5 The equation of the tangent to the circle at K in the form $y = mx + c$ (2)

4.2* Determine the values of t such that the line $y = \frac{1}{2}x + t$ cuts the circle at two different points. (3)

4.3* Calculate the area of $\triangle SMK$. (5) [23]

See pp. xi & xii for Analytical Geometry Toolkit



▶ TRIGONOMETRY [41]


QUESTION 5 Answers on p. A18

Trig Summary on p. vii



5.1 Given: $\frac{\sin(A - 360^\circ) \cdot \cos(90^\circ + A)}{\cos(90^\circ - A) \cdot \tan(-A)}$
Simplify the expression to a single trigonometric ratio. (6)

5.2 In the diagram, $P(3; t)$ is a point in the Cartesian plane. $OP = \sqrt{34}$ and reflex $\hat{HOP} = \beta$.



Without using a calculator, determine the value of:


- 5.2.1 t 5.2.2 $\tan \beta$ 5.2.3 $\cos 2\beta$ (2)(1)(4)

5.3* Prove:

- 5.3.1 $\sin(A + B) - \sin(A - B) = 2 \cos A \cdot \sin B$ (2)
- 5.3.2 and hence, without using a calculator, that $\sin 77^\circ - \sin 43^\circ = \sin 17^\circ$ (4) [19]

QUESTION 6 Answers on p. A19

In the diagram, the graph of $f(x) = \cos 2x$ is drawn for the interval $x \in [-270^\circ; 90^\circ]$.



6.1 Draw the graph of $g(x) = 2 \sin x - 1$ for the interval $x \in [-270^\circ; 90^\circ]$ on the grid. Show ALL the intercepts with the axes, as well as the turning points. (4)


6.2* Let A be a point of intersection of the graphs of f and g . Show that the x -coordinate of A satisfies the equation $\sin x = \frac{-1 + \sqrt{5}}{2}$. (4)

6.3* Hence, calculate the coordinates of the points of intersection of graphs of f and g for the interval $x \in [-270^\circ; 90^\circ]$. (4) [12]

QUESTION 7 Answers on p. A20

AB represents a vertical netball pole. Two players are positioned on either side of the netball pole at points D and E such that D, B and E are on the same straight line. A third player is positioned at C . The points B, C, D and E are in the same horizontal plane.

The angles of elevation from C to A and from E to A are x and y respectively. The distance from B to E is k .



7.1 Write down the size of \hat{ABC} . (1)

7.2* Show that $AC = \frac{k \cdot \tan y}{\sin x}$ (4)

7.3* If it is further given that $\hat{DAC} = 2x$ and $AD = AC$, show that the distance DC between the players at D and C is $2k \tan y$. (5) [10]

Consult the Topic Guide on page 2 for more examples on a particular topic.



2.3 All the terms from -3 onwards are negative

∴ There are 22 terms, i.e. $n = 22$

$a = -3$; $d = -4$; $S_{22}?$; $T_{22} = -87$

$S_n = \frac{n}{2}(a + T_n) \Rightarrow S_{22} = \frac{22}{2}[-3 + (-87)]$

$= 11(-90)$

$= -990 <$

Always preferable to use this formula when you have the value of T_n .

OR: $S_n = \frac{n}{2}[2a + (n - 1)d]$

$\Rightarrow S_{22} = \frac{22}{2}[2(-3) + (22 - 1)(-4)]$

$= 11[-6 - 84]$

$= 11[-90]$

$= -990 <$

2.4 Now, $a = 5$ & $d = -4$, but $T_n = -4 187$, $n?$

$T_n = -4n + 9 \Rightarrow -4 187 = -4n + 9$

$\therefore -4 196 = -4n$

$\therefore n = 1 049$

OR: $T_n = a + (n - 1)d \Rightarrow -4 187$

$= 5 + (n - 1)(-4)$, etc.

Terms divisible by 5:


5 ; 1 ; -3 ; -7 ; -11 ; -15 ; -19 ; -23 ; -27 ; -31 ; -35 ; ...

The first term of each group of 5 terms is divisible by 5.

The number of groups of 5 terms = $\frac{1 049}{5} = 209 \text{ rem } 4$.

∴ There are 209 complete groups and the first 4 terms of the next group (which will start with a multiple of 5).

∴ 210 terms <



3.1.1 $2 - 2x = 2 + 2x$... quadratic sequence has equal 2nd differences
 $\therefore -4x = 0$
 $\therefore x = 0 <$



The 1st differences: 1 ; 3 ; 5 ; ...

$\therefore T_n = 2n - 1$

& $S_n = \frac{n}{2}(a + T_n)$

$= \frac{n}{2}(1 + 2n - 1)$

$= \frac{n}{2}(2n)$

$= n^2$

$S_n = 250 \Rightarrow n^2 = 250$

$\therefore n = 15,81... \dots n > 0$

$\therefore S_n > 250 \Rightarrow n = 16$

The 16th term (of the 1st differences) is the 1st difference between the 16th and 17th term of the original number pattern

∴ The required position is the 17th term <

Note: The positions of the first differences are 1 behind the positions of the original number pattern.

3.2.1 The lengths of the rectangles are:

21 ; $21 \times 0,85$; $21 \times (0,85)^2$; ...

$\therefore T_{10} = 21 \times 0,85^9 \dots T_n = ar^{n-1}$

$\approx 4,86 \text{ cm} <$

3.2.2 (The length of the sheet is 30 cm
 \therefore 15 rectangles can be drawn)

The sum of the areas of the 15 rectangles:

S_{15}

$= 21 \times 1 + 21 \times 0,85 \times 1 + 21 \times (0,85)^2 \times 1 + \dots \dots 15 \text{ terms}$

$= \frac{21 [1 - (0,85)^{15}]}{1 - 0,85} \dots S_n = \frac{a(1-r^n)}{1-r}$

$= 127,77 \text{ cm}^2$

\therefore The % that is shaded = $\frac{127,77}{21 \times 30} = 0,20281...$

$\approx 20,28\% <$

► FUNCTIONS AND GRAPHS [35]

4.1 $y = 0 <$... The asymptote is the x-axis!
 & The eqn. of the x-axis is $y = 0$

4.2 $R(0; 1) <$... At R, $x = 0$ & $y = a^0 = 1$

4.3 Pt P(2; 9) on graph $y = a^x$
 Substitute: $\therefore 9 = a^2$
 $\therefore a = 3 <$... $a \geq 0$ in $y = a^x$

4.4 Pt Q($b; \frac{1}{81}$) on graph $y = 3^x$... $a = 3$ in 4.3
 $\therefore \frac{1}{81} = 3^b$
 $\therefore 3^b = 3^{-4} \dots \frac{1}{81} = \frac{1}{3^4} = 3^{-4}$
 $\therefore b = -4$

$\therefore x_D = -4 \dots x_D = x_Q$

\therefore Length of DP = $4 + 2 \dots$ or, $2 - (-4)$
 $= 6 \text{ units} <$



8.1.5 In $\triangle ONR$:


$$\begin{aligned} ON &= OR \quad \dots \text{equal radii} \\ \therefore \hat{ONR} &= \hat{R} \\ &= \frac{1}{2}(180^\circ - 66^\circ) \quad \dots \text{sum of } \angle^s \text{ of } \triangle \\ &= 57^\circ \\ \therefore \hat{N}_2 &= 57^\circ - 24^\circ \\ &= 33^\circ \end{aligned}$$



8.2.1 $FC \parallel AB$... opp. sides of a rectangle

& $AB \parallel GH$... given

$\therefore FC \parallel GH$ <



8.2.2 In $\triangle AGH$:

$$\frac{AC}{CH} = \frac{AF}{FG} \quad \dots \text{prop. theorem; } FC \parallel GH$$

$$\therefore \frac{AC}{21} = \frac{20}{15}$$

$$\therefore AC = \frac{21 \times 20}{15} = 28 \text{ units}$$

$\therefore DB (= AC) = 28$ units ... Diagonals of a rect. are =

$$\therefore DM = \frac{1}{2}(28) \quad \dots \text{Diagonals of a } \square \text{ (and } \therefore \text{ a rectangle) bisect one another}$$


$$= 14 \text{ units} <$$

9.1 Theorem

$$\text{RTP: } \hat{J} + \hat{L} = 180^\circ$$

Construction:

Draw radii OK and OM



Proof:


$$\hat{J}_1 = \frac{1}{2}\hat{O}_1 \text{ and } \hat{L} = \frac{1}{2}\hat{O}_2 \quad \dots \angle \text{ at centre} = 2 \times \angle \text{ at circumference}$$

$$\therefore \hat{J}_1 + \hat{L} = \frac{1}{2}(\hat{O}_1 + \hat{O}_2)$$

But $\hat{O}_1 + \hat{O}_2 = 360^\circ$... \angle^s about point O

$$\therefore \hat{J}_1 + \hat{L} = 180^\circ <$$

9.2



9.2.1 (a) $\hat{B}_1 = x$ < ... \angle^s in the same segment; chord ST subtends

(b) $\hat{B}_2 = y$ < ... exterior \angle of cyclic quadrilateral

9.2.2 Using 9.2.1 (a) & (b): $\hat{SBD} = x + y$

& In $\triangle ACR$: $\hat{C} = 180^\circ - (x + y)$... sum of \angle^s of \triangle

\therefore In quadrilateral SCDB:

\hat{SBD} and \hat{C} are supp. \angle^s

\therefore SCDB is a cyclic quad. <

... CONVERSE of 'opp. \angle^s of cyclic quad.' theorem

Here, we use the CONVERSE of the 'opp. \angle^s of c.q.' thm.

9.2.3

We need to prove: $x + y \neq 90^\circ$

$$\begin{aligned} \hat{C} &= \hat{AST} - \hat{D}_2 \quad \dots \text{ext. } \angle \text{ of } \triangle SCD \\ &= 100^\circ - 30^\circ \\ &= 70^\circ \end{aligned}$$

$\therefore \hat{SBD} (= x + y) = 110^\circ$... opp. \angle^s of c.q. SCDB

$\therefore x + y \neq 90^\circ$

\therefore SD is not a diameter of circle BDS <

... CONVERSE of ' \angle in semi- \odot ' thm.




10.1.1 We need to prove that

$$\hat{C}_2 = \hat{A}_1$$



We will use the CONVERSE of the tan chord theorem



$$\begin{aligned} \hat{D}_2 &= \hat{B} \quad \dots \text{exterior } \angle \\ &= x \quad \dots \text{of cyclic quad.} \end{aligned}$$

\therefore In $\triangle DMC$:

$$\hat{C}_2 = 90^\circ - x \quad \dots \text{sum of } \angle^s \text{ of } \triangle$$

But $\hat{A}_1 = \hat{A}_2$... equal chords subtend equal angles

& In $\triangle ACB$: $\hat{A}_2 = 90^\circ - x$... sum of \angle^s of \triangle

$$\therefore \hat{A}_1 = 90^\circ - x$$

$$\therefore \hat{C}_2 = \hat{A}_1$$

\therefore MC is a tangent to the circle at C <

... CONVERSE of tan chord theorem

10.1.2 In \triangle^s ACB and CMD

(1) $\hat{ACB} = \hat{M} (= 90^\circ)$... given

(2) $\hat{A}_2 = \hat{C}_2$... both $= 90^\circ - x$ in 10.1.1

$\therefore \triangle ACB \parallel \triangle CMD$ < ... $\angle \angle \angle$

$$10.2.1 \frac{CM}{DC} = \frac{AC}{AB} \quad \dots \textcircled{1} \quad \dots \text{similar } \triangle^s \text{ in 10.1.2}$$

But, in \triangle^s CMD and AMC

(1) $\hat{M} (= 90^\circ)$ is common

(2) $\hat{C}_2 = \hat{A}_1$... proved in 10.1.1

$\therefore \triangle CMD \parallel \triangle AMC$... $\angle \angle \angle$

$$\therefore \frac{CM}{DC} = \frac{AM}{AC} \quad \dots \textcircled{2} \quad \dots \parallel \triangle^s$$

$$\therefore \frac{CM}{DC} \times \frac{CM}{DC} = \frac{AC}{AB} \times \frac{AM}{AC} \quad \dots \text{see } \textcircled{1} \text{ and } \textcircled{2}$$

$$\therefore \frac{CM^2}{DC^2} = \frac{AM}{AB} <$$

$$10.2.2 \frac{AM}{AB} = \frac{CM^2}{DC^2} \quad \dots \text{proved in 10.2.1}$$

But, in $\triangle DMC$: $\frac{CM}{DC} = \sin x$

$$\therefore \frac{AM}{AB} = \sin^2 x <$$




► The Proportion Theorem

6

A line parallel to one side of a triangle divides the other two sides proportionally.


i.e. $DE \parallel BC \Rightarrow \frac{AD}{DB} = \frac{AE}{EC}$



Given: $\triangle ABC$ with $DE \parallel BC$,
D & E on AB & AC respectively.

To prove: $\frac{AD}{DB} = \frac{AE}{EC}$

Construction: Join DC & BE



Proof: $\frac{\text{Area of } \triangle ADE}{\text{Area of } \triangle DBE} = \frac{\frac{1}{2}AD \cdot h}{\frac{1}{2}DB \cdot h} = \frac{AD}{DB}$

h is the height of $\triangle^s ADE$ and DBE



Similarly: $\frac{\text{Area of } \triangle ADE}{\text{Area of } \triangle EDC} = \frac{\frac{1}{2}AE \cdot h'}{\frac{1}{2}EC \cdot h'}$

h' is the height of $\triangle^s ADE$ and EDC

But: $\triangle DBE = \triangle EDC$, in area ... *on the same base DE ; between || lines, DE & BC*

and: $\triangle ADE$ is common

$\therefore \frac{\text{Area of } \triangle ADE}{\text{Area of } \triangle DBE} = \frac{\text{Area of } \triangle ADE}{\text{Area of } \triangle EDC}$


$\therefore \frac{AD}{DB} = \frac{AE}{EC} \leftarrow$



► The Similar \triangle^s Theorem

7

If two triangles are equiangular, then their sides are proportional and, therefore, they are similar.



Given: $\triangle ABC$ & $\triangle DEF$ with $\hat{A} = \hat{D}$ $\hat{B} = \hat{E}$ & $\hat{C} = \hat{F}$

To prove: $\frac{AB}{DE} = \frac{AC}{DF} = \frac{BC}{EF}$

Construction: Mark P & Q on DE & DF such that DP = AB & DQ = AC

Proof: In $\triangle^s DPQ$ & ABC

(1) DP = AB ... construction

(2) DQ = AC ... construction

(3) $\hat{D} = \hat{A}$... given

$\therefore \triangle DPQ \equiv \triangle ABC$... $S\angle S$

$\therefore \hat{P}_1 = \hat{B}$
 $= \hat{E}$... given

$\therefore PQ \parallel EF$... corresponding \angle^s equal

$\therefore \frac{DP}{DE} = \frac{DQ}{DF}$... proportion theorem;
 $PQ \parallel EF$

But DP = AB and
DQ = AC ... construction

$\therefore \frac{AB}{DE} = \frac{AC}{DF}$

Similarly, by marking S and T on DE and EF such that

$SE = AB$ and $ET = BC$, it can be proved that: $\frac{AB}{DE} = \frac{BC}{EF}$

$\therefore \frac{AB}{DE} = \frac{AC}{DF} = \frac{BC}{EF} \leftarrow$

$\therefore \triangle ABC$ and $\triangle DEF$ are similar.



The focal point

1 congruency

2 corresponding \angle^s

3 parallel lines

4 proportions



Similar \triangle^s



\triangle^s are similar if: **A:** they are equiangular, and
B: their sides are proportional

In this proof, we show that: **A** \rightarrow **B**
i.e. Both conditions, **A** and **B**, apply

\therefore The \triangle^s are similar

The converse statement says: **B** \rightarrow **A**

i.e. Both conditions, **A** and **B**, apply

\therefore The \triangle^s are similar

Compound Angle Formulae



1. $\sin(A + B) = \sin A \cos B + \cos A \sin B$ *Sign stays the same sine & cosine of A and B mixed*
2. $\sin(A - B) = \sin A \cos B - \cos A \sin B$
3. $\cos(A + B) = \cos A \cos B - \sin A \sin B$ *Sign changes cosine of A and B first, then sine of A & B*
4. $\cos(A - B) = \cos A \cos B + \sin A \sin B$

We will prove formula no. 4 (see above) and then derive the other 3 from it.



Double Angle Formulae



5. $\sin 2A = 2 \sin A \cos A$... This formula will be derived from the formula no. 1.
6. $\cos 2A = \cos^2 A - \sin^2 A$... This formula will be derived from the formula no. 3.

or $\cos 2A = 1 - 2 \sin^2 A$


or $\cos 2A = 2 \cos^2 A - 1$



Proof of the Formula:

$$\cos(A - B) = \cos A \cos B + \sin A \sin B$$

First, an important concept!



NOTE: If $OP = 1$ unit!

then: $\frac{x}{1} = \cos \theta$ and $\frac{y}{1} = \sin \theta$

i.e. $x = \cos \theta$ and $y = \sin \theta$

i.e. **P is the point $(\cos \theta; \sin \theta)$**


In the sketch alongside, \hat{A} and \hat{B} have been placed in standard position.

$$\hat{R}\hat{O}\hat{Q} = \hat{A} - \hat{B}$$

The coordinates of the points **R** and **Q**, both **1 unit** from the origin, are:

R $(\cos A; \sin A)$ & **Q** $(\cos B; \sin B)$

... See NOTE above



► Determine 2 expressions for RQ^2

$$RQ^2 = 1^2 + 1^2 - 2(1)(1) \cos(A - B) \quad \dots \quad \text{COSINE RULE}$$

$$= 2 - 2 \cos(A - B) \quad \dots \quad \textcircled{1}$$

& $RQ^2 = (\cos A - \cos B)^2 + (\sin A - \sin B)^2 \quad \dots \quad \text{DISTANCE FORMULA}$

$$= \cos^2 A - 2 \cos A \cos B + \cos^2 B + \sin^2 A - 2 \sin A \sin B + \sin^2 B$$

$$= 2 - 2 \cos A \cos B - 2 \sin A \sin B \quad \dots \quad \textcircled{2} \quad \dots \quad \sin^2 \theta + \cos^2 \theta = 1$$

► Equate the two expressions for RQ^2 above:

$$\textcircled{1} = \textcircled{2} \quad \therefore 2 - 2 \cos(A - B) = 2 - 2 \cos A \cos B - 2 \sin A \sin B$$


► Subtract 2: $\therefore -2 \cos(A - B) = -2 \cos A \cos B - 2 \sin A \sin B$

► Divide by -2 (or \times by $-\frac{1}{2}$): $\therefore \cos(A - B) = \cos A \cos B + \sin A \sin B <$

QUADRILATERALS - definitions, areas & properties

All you need to know

'Any' Quadrilateral



Sum of the \angle^s of any quadrilateral = 360°


$$\begin{aligned} \text{Sum of the interior angles} &= (a + b + c) + (d + e + f) \\ &= 2 \times 180^\circ \dots (2\Delta^s) \\ &= 360^\circ \end{aligned}$$

The arrows indicate various 'pathways' from 'any' quadrilateral to the square (the 'ultimate quadrilateral'). These pathways, which combine logic and fact, are essential to use when proving specific types of quadrilaterals.

See how the properties accumulate as we move from left to right, i.e. the first quad. has no special properties and each successive quadrilateral has all preceding properties.



A Trapezium




DEFINITION:
Quadrilateral with 1 pair of opposite sides \parallel

$$\begin{aligned} \text{Area} &= \Delta 1 + \Delta 2 \\ &= \frac{1}{2} ah + \frac{1}{2} bh \\ &= \frac{1}{2} (a + b) \cdot h \end{aligned}$$


'Half the sum of the \parallel sides \times the distance between them.'

A Parallelogram



DEFINITION:
Quadrilateral with 2 pairs opposite sides \parallel

Area = base \times height




\parallel^m ABCD = Δ BCQ + Δ QCD
rect. PBCQ = Δ BCQ + Δ PBA
where Δ QCD \cong Δ PBA \dots RHS/ 90° HS
 $\therefore \parallel^m$ ABCD = rect. PBCQ (in area)
= BC \times QC

Properties:

- 2 pairs opposite sides equal
- 2 pairs opposite angles equal
- & **DIAGONALS BISECT ONE ANOTHER**

A Rectangle



DEFINITION:
A \parallel^m with one right \angle

$$\text{Area} = \ell \times b$$

DIAGONALS are EQUAL

The Square






the 'ultimate' quadrilateral!


$$\text{Area} = s^2$$

Properties:

It's all been said 'before'!
Since a square is a rectangle, a rhombus, a parallelogram, a kite, \dots ALL the properties of these quadrilaterals apply.



A Kite



DEFINITION:
Quadrilateral with 2 pairs of adjacent sides equal

Given diagonals a and b


$$\text{Area} = 2\Delta^s = 2 \left(\frac{1}{2} b \cdot \frac{a}{2} \right) = \frac{ab}{2}$$

'Half the product of the diagonals'

THE DIAGONALS

- cut perpendicularly
- ONE DIAGONAL** bisects the other diagonal, the opposite angles and the area of the kite

A Rhombus



DEFINITION:
A \parallel^m with one pair of adjacent sides equal

Area

$$= \frac{1}{2} \text{product of diagonals (as for a kite)}$$


or

$$= \text{base} \times \text{height (as for a parallelogram)}$$

THE DIAGONALS

- bisect one another **PERPENDICULARLY**
- bisect the angles of the rhombus
- bisect the area of the rhombus

Note:






$$2x + 2y = 180^\circ \dots \angle^s \text{ of } \Delta \text{ or}$$

$$\rightarrow x + y = 90^\circ \text{ co-int. } \angle^s; \parallel \text{ lines}$$



CONCAVITY & THE POINT OF INFLECTION

The **Concavity** of cubic graphs: **Concave up**  or **Concave down** , changes at the point of inflection:
As x increases (i.e. from left to right) ...




Note: For cubic graphs with 2 stationary points, the coordinates of the point of inflection are the averages of the x - and y -coordinates of the stationary points.

$f''(x) = 0$ at the point of inflection of f
 $f''(x) < 0$ where f is concave down
 $f''(x) > 0$ where f is concave up

GROUPING OF CIRCLE GEOMETRY THEOREMS

The grey arrows indicate how various theorems are used to prove subsequent ones



There are '2 ways' to prove that a line is a tangent to a \odot !